

The Effectiveness of Acceptance and Commitment Therapy on Externalizing Symptoms in 7 to 12 Year-Old Children with Chronic Pain

Original Article

Soheila Ghomian (PhD) *¹
Mohammad Reza Shairi (PhD) ²

1. Faculty of Humanities, Shahed University, Tehran, IR Iran.
2. Associate Professor of Clinical Psychology, Faculty of Humanities, Shahed University, Tehran, IR Iran.

* Correspondence:

Soheila Ghomian, Student in Clinical Psychology. Faculty of Humanities, Shahed University, Persian Gulf Highway, 33191-18651, Tehran, Iran.

E-mail:

soheila_ghomian@yahoo.com

Tel: +98 2151212470

Fax: +98 2151213564

Received: 9 Dec 2015

Revised: 6 Jan 2016

Accepted: 1 Feb 2016

Abstract:

Background: In recent years, the interest in relation to the identification of adaptive mechanisms through which people continue to improve their psychological well-being, despite the experience of chronic pain, has been observed. Acceptance is one of these positive psychological factors. We aimed to investigate the effect of Acceptance and Commitment Therapy for Children with Chronic Pain on externalizing symptoms in 7 to 12 year-old children.

Methods: The present study was based on quasi-experimental model According to the criteria of chronic pain; a number of children with chronic pain were selected by available sampling method from specialty and subspecialty pediatric hospitals of Tehran. Then, 20 children who according to their parents were prepared to participate in this study and met the inclusion criteria, were selected and were placed in the experimental and control group. The Child Behavior Checklist was administered in both groups at the pre-test, post-test, first and second follow-up.

Results: The results showed that the experimental group compared with the control group showed significant change in externalizing symptoms in multiple stages. These changes continued after the treatment, primary and secondary follow-up. Chi-square related to total score in experimental group was 26.21 ($p=0.001$) and chi-square related to total score in control group was 18.55 ($p=0.069$).

Conclusions: Generally, it can be said that this protocol can be used in clinical fields, especially in the area of improving the externalizing symptoms of children.

Keywords: Acceptance and Commitment Therapy, Chronic Pain, Externalizing Symptoms

Citation:

Ghomian S, Shairi MR. The Effectiveness of Acceptance and Commitment Therapy on Externalizing Symptoms in 7 to 12 Year-Old Children with Chronic Pain. Caspian J of Pediatr March 2016; 2(1): 107-12.

Introduction:

Emotive regulation feature is an adaptive solution, especially when dealing with anger and distress. This feature leads to increased psychological well-being ^[1]. Emotive regulation is the process which individuals automatically or consciously reduce their emotions ^[2,3] and try to respond effectively to the demands of the environment with the influence on the occurrence, severity, duration, and expression of these emotions ^[3,4]. Although the current evidence suggests that emotive dysregulation is effective on the formation and persistence of emotional disorders, for example externalizing disorders (e.g. aggressive and offensive behavior) ^[5-7], growing evidence indicates that this concept has found wide application in the domains of pain and physical health. Preliminary findings have emphasized the important role of emotive regulation in painful reactions, especially from the angle of the problems associated with identifying and expressing emotions.

The research of Van Middendorp et al.'s showed that in patients with chronic pain, who were faced with the problem of identifying and expressing emotions, severe emotion was associated with more severe pain. These findings indicated that interventions aimed at improving emotive regulation and psychological flexibility (in response to different emotional situations), might facilitate the chronic pain improvement [8].

On the other hand, because among the different ways of emotive regulation, acceptance is lower among individuals with externalizing symptoms [9] and the results of some studies indicate the impact of acceptance strategy on emotive regulation [10-12], therefore, it can be concluded that interventions that are aimed at increasing acceptance and psychological flexibility have striking effects on externalizing symptoms in children with chronic pain.

Acceptance and commitment therapy (ACT) is an acceptance-based therapy that has empirical support for many physical and emotional problems, and research evidence for ACT of chronic pain shows an increasing trend [13].

ACT includes the non-judgmental awareness and acceptance of all experiences (positive and negative experiences), identification of valuable orientation of life, and efficient operation based on the goals that are consistent with these valuations. The target of ACT is clearly improving the performance and quality of life through the experience of thoughts, emotions and negative feelings of body with a more flexible manner. The process of ACT is a cycle of detecting cognitive fusion and avoidance, defusing and letting go (thus establishing new, more flexible functions for these events) and moving in a valued direction in a way that builds larger and larger patterns of effective behavior [14].

Various interventions have been conducted in relation to reducing the externalizing symptoms of children. For example, the research of Lochman et al.'s has demonstrated the effectiveness of cognitive behavior therapy (CBT: is a form of psychotherapy that change unhelpful thinking and behavior) strategies on externalizing symptoms [15].

Mindfulness therapy (mindfulness is the gentle effort to be continuously present with experience) is recognized as another effective intervention on externalizing symptoms. For example, the research of Bogels et al.'s showed that after mindfulness training, the children gained a better improvement in individual goals, internalizing problems, externalizing problems,

attention problems and happiness. Moreover, they showed better performance in sustained attention tests and parents reported greater improvement in their children's goals, internalizing problems, externalizing problems, and their self-control [16].

Although there are various interventions in reducing children's externalizing symptoms, there is a research gap in the area of reducing externalizing symptoms in children with chronic pain, especially based on ACT. Thus, on one hand, the existing research gap in this area and on the other hand, because of the negative consequences of externalizing behavior [17], such as social and educational problems [18], the basic problem of the current study was whether Acceptance and Commitment Therapy for Children with Chronic Pain (CHACT) [19] (CHACT was similar to ACT, but was designed to fit children) could improve the externalizing symptoms of 7 to 12 year-old children with chronic pain or not?

Methods:

The present study was based on quasi-experimental model. Among the patients referred to different clinics and departments of specialty and subspecialty pediatric hospitals in Tehran such as Mofid Children's Hospital, Children's Medical Center, Hazrat Ali Asghar Hospital and Bahrami Hospital (In these centers, different parts and clinics were used, such as: neurology, neurosurgery, surgery, blood, rheumatology, orthopedics and physiotherapy), 20 children who met the inclusion criteria were entered into the study according to their parents' consent. Sampling method of this study was based on available sampling method.

Inclusion criteria for this study were as follows: (1) Being in the age range of 7 to 12 years old, (2) Engagement with education (because, children who are not in education employment, may have difficulty in understanding treatment and assignments); evaluation of educational status (success or failure), according to the school status, was done by the psychologist, (3) Obtaining the score of 13 to 29 (moderate disability) in Function Disability Inventory (FDI) (children who their score was less than 13, chronic pain did not have effect on their performance), (4) Having a history of developing chronic pain for 6 months or more and at least 3 months of the first medical treatment in relation to chronic pain and (5) The ability to attend meetings. After the selection of subjects based on inclusion criteria, they were placed in the experimental group (n=10) and control group (n=10).

Then, CHACT was implemented on the experimental group. This protocol was designed based on the books of ACT, initial grete of ACT on children, ACT on adults with chronic pain, model of anxiety treatment in children, model of OCD treatment in children and consultation with Association for Contextual Behavioral Science (ACBS) (such as doctor Hayes, Wicksell, Murrell and Wilson). CHACT was based on six components related to ACT, such as acceptance, diffusion, and contact with the present moment, values, commitment action and self as context. More details about the protocol, such as templates and the content of the meetings were given in the previous paper [19].

Tools: The used tools in this study were as follows:

1) Demographic questionnaire:

The questions were about age, sex, education, chronic pain criteria (a history of developing chronic pain for 6 months or more, according to the approved physician and at least 3 months of the first medical treatment in relation to the chronic pain), taking or not taking pain medication, type and amount of pain medication (if used), and education and occupation of parents.

2) Child Behavior Checklist (CBCL)

CBCL assesses competencies (abilities), action or adaptive functioning and emotional-behavioral problems of 6-18 year-old children. This checklist which contained 13 questions and 113 items was answered by parents or a person who was responsible for the child and took care of him/her. The CBCL scales included competence and adaptive functioning scales, scales based on experience and scales based on DSM. The raw scores of problem CBCL scales were obtained by summing the scores of 0, 1 or 2 questions related to any scale. Minaee (2006) obtained good psychometric properties of CBCL [20]. In this study, we used the scales based on experience.

CBCL was used before and after the treatment and first follow-up (1.5 months after the treatment) and second follow-up (5 months after the treatment).

Method of data analysis: In this study besides descriptive statistics, because of the lack of the assumptions related to parametric tests, Friedman Test was used for examination of change in different time periods and Mann-Whitney Test was used for comparison of difference between the groups in the pre-test, post-test, first and second follow-up. SPSS19 was used for data analysis. Statistical analysis was performed in Shahed University. (IRCT No.: 2013112815577).

Results:

The results are presented in two sections of descriptive and analytical results:

A) Descriptive results:

The descriptive results of this study suggested that the mean age (SD) of the experimental and control group was 10.60 (1.7) and 10.20 (1.81), respectively, and the youngest children were 7 years old. The experimental group consisted of 4 girls and 6 boys and the control group consisted of 5 girls and 5 boys. In both groups, most patients were suffering from chronic pain caused by rheumatoid disease and the rest was suffering from pain in the chest, leg, kidney and so on. Many subjects in both groups were taking medication (11 person). In both groups, many parents were educated in guidance school.

Table1 presents the descriptive indicators of externalizing symptoms variable and its subscales. As can be seen, in the experimental group, external and aggressive behavior subscales have changed from pretest (M (SD)=6 (3.74) M (SD)=7.90, respectively) to posttest (M (SD)=2.70 (2) M (SD)=4.10 (2.80), respectively) and have remained relatively constant in the first and second follow-up. In the control group, externalizing behavior subscales remained relatively constant in all four time sections. Significant and non-significant statistical results of this status will be presented in the next section.

B) Analytical results:

Before addressing these results, it is worth mentioning that in both groups, based on parents' views, comparing the externalizing behavior variable in pre-test base on Mann-Whitney Test is not significant. (Offensive behavior, -1.013 (0.311); aggressive behavior, -0.228 (0.820); total score, -0.227 (0.820)). The analytical results of this study are presented in Tables 2 and 3. As shown in Table 2, in experimental groups, externalizing and aggressive behavior subscales are obtained significant changes at different time sections (chi-square for aggressive behavior = 23.28 (p=0.001), chi-square for externalize behavior (total)=26.21 (p=0.001)). Table 3 shows the meaningful comparison of variables between the control and experimental groups. As illustrated in this table, both groups generally show significant differences in relation to many variables. The most difference between the two groups was seen at this time: between pretest with posttest (z (total)=-2.28 (p=0.020)), between pretest with follow up 1 (z (total)=-2.19 (p=0.023)) and between pretest with follow up 2 (z (total)=-2.08 (p=0.021)). Thus, in

general, we can say that the CHACT is able to improve the externalizing symptoms of 7 to 12 year-old children

with chronic pain.

Table 1- Mean (SD) of the studied variables in experimental and control groups based on the responses of parents

| Variable | Groups | Pretest | Posttest | Follow up 1 | Follow up 2 |
|--------------------------------|--------------|-------------|-------------|-------------|-------------|
| Aggressive behavior | Experimental | 6 (3.74) | 2.70 (2) | 2.80 (2.20) | 2.60 (2.30) |
| | Control | 6.10 (5.40) | 5 (4.54) | 5.10 (4.60) | 4.30 (4.70) |
| Offense behavior | Experimental | 1.90 (1.37) | 1.40 (1.07) | 1.50 (1.17) | 1.43 (1.10) |
| | Control | 1.50 (1.84) | 1.10 (1.59) | 1.30 (1.70) | 1.22 (1.50) |
| Total (externalizing behavior) | Experimental | 7.90 (4.84) | 4.10 (2.80) | 4.30 (3.09) | 3.90 (2.80) |
| | Control | 7.60 (6.27) | 6.10 (5.23) | 6.20 (5.26) | 5.90 (5.03) |

Table 2- Externalizing symptoms change and its subscales in the four time; pre-test, post-test, first and second follow-up

| Groups | Variable | Chi - Square | df | P value |
|--------------|--------------------------------|--------------|----|---------|
| Experimental | Aggressive behavior | 23.286 | 3 | .001** |
| | Offense behavior | 6.000 | 3 | .112 |
| | Total (externalizing behavior) | 26.217 | 3 | .001** |
| Control | Aggressive behavior | .360 | 3 | .948 |
| | Offense behavior | 9.000 | 3 | .054 |
| | Total (externalizing behavior) | 18.558 | 3 | .069 |

*P<0.05 **P<0.01

Table 3- Comparison of externalizing symptoms in experimental and control group

| | | Aggressive behavior | | Offense behavior | | Total (externalizing behavior) | |
|------------------------|-------------|---------------------|--------------|------------------|-------------|--------------------------------|--------------|
| | | Experimental g | Control g | Experiment | Control | Experiment | Control |
| Pretest & post test | Mean (SD) | 3.30 (2.49) | 1.10 (3.75) | 0.50 (0.84) | 0.40 (0.69) | 3.80 (2.78) | 1.50(4.30) |
| | Z (P value) | -2.148 (0.023)* | | -0.140 (0.912) | | -2.285 (0.020)* | |
| Pretest & follow up 1 | Mean (SD) | 3.20 (2.61) | 1.00 (3.77) | 0.40 (0.96) | 0.38 (0.70) | 3.60 (2.95) | 1.40 (4.32) |
| | Z (P value) | -2.326 (0.019)* | | -0.178 (0.812) | | -2.196 (0.023)* | |
| Pretest & follow up 2 | Mean (SD) | 3.29 (2.49) | 1.10 (3.70) | 0.48 (0.88) | 0.40 (0.67) | 3.72 (2.82) | 1.54 (4.20) |
| | Z (P value) | -2.200 (0.013)* | | -0.130 (0.712) | | -2.085 (0.021)* | |
| Posttest & follow up 1 | Mean (SD) | -0.10 (0.56) | -0.13 (0.31) | -0.10 (0.31) | 0.00 (0.00) | -0.20 (0.78) | -0.10 (0.31) |
| | Z (P value) | -0.054 (0.971) | | -1.000 (0.739) | | -0.108 (0.871) | |
| Posttest & follow up 2 | Mean (SD) | 0.00 (0.00) | 0.00 (0.00) | 0.00 (0.00) | 0.00 (0.00) | 0.00 (0.00) | 0.00 (0.00) |
| | Z (P value) | 0.000 (1.000) | | 0.000 (1.000) | | 0.000 (1.000) | |
| Follow up 1 & 2 | Mean (SD) | 0.09 (0.56) | 0.10 (0.31) | 0.15 (0.28) | 0.00 (0.00) | 0.20 (0.70) | 0.10 (0.31) |
| | Z (P value) | -0.054 (0.961) | | -1.004 (0.639) | | -0.128 (0.971) | |

received CHACT compared to children who did not receive this treatment.

The results of previous research have shown that anger management, social problem solving and goal setting are the effective components in the treatment of children with externalizing symptoms and in this respect, the effectiveness of CBT for reducing the externalizing symptoms of children has been proven

Discussion:

The purpose of this study was to investigate the effect of CHACT on externalizing symptoms of 7 to 12 year-old children with chronic pain. Overall, this research indicated that there were significant changes in externalizing symptoms among children who

[15]. Thus, It can be concluded that in this study, the components of acceptance, cognitive fusion (according to its purpose in relation to the reduction in the true sense of inner experiences and also, attention to thoughts just as thought), valuing (determining objectives and values associated with them) and committed action (according to the goal that has in relation to skills, achieving values and the creation of larger patterns of effective and flexible behavior), are the most influential factors in reducing children's externalizing symptoms. As indicated, among the subscales of externalizing symptoms, offense behavior subscale showed no significant change in the experimental group comparing to the control group. It seems that one of the reasons is the role of different variables influencing the behavior of rule violations, for example the negative stereotypes of parents [21] and subsequently, more anger of parents towards these behavior [22], high levels of criticism [23] and observed conflicts in these families [21], parenting style and parental support [21], similar or different parenting styles of mother and father [25], low self-esteem of children with offensive behavior [26] and the importance of the school environment [27].

According to the research gaps related to the effect of ACT on externalizing symptoms in children with chronic pain, we can say that generally the present results which are consistent with the results of Lochman et al.'s and Bogels et al.'s respectively, have proved the efficacy of CBT and mindfulness on externalizing symptoms in children [15,16].

Regarding the impact of CHACT on externalizing symptoms of the children with chronic pain, it can be said that this protocol can be used in clinical fields, especially in the area of improving the externalizing symptoms of children. Although this study was conducted to determine the impact of CHACT on children with chronic pain regarding the limited sample size, it required multiple checks in terms of economic, social and cultural diversity in Iran. There is no doubt that if the conditions of comparing CHACT with other psychological treatment of chronic pain in children and also with the "placebo" were obtained, its role and impact would be more seriously evaluated.

Acknowledgment:

We would like to extend our deep appreciations to the staff of the hospitals involved in this study, particularly the presidency section that let us do this research. In this study, corresponding author was

involved in the all process (such as sampling, treatment and so on) and the second author, as a supervisor, indicated general guidance.

Funding: This study was self-funded.

Conflict of interest: There was no conflict of interest.

References:

1. Mauss IB, Cook CL, Cheng JY, Gross JJ. Individual differences in cognitive reappraisal: experiential and physiological responses to an anger provocation. *International journal of psychophysiology: offic J Int Organiz Psychophysiol* 2007; 66(2): 116-24.
2. Bargh JA, Williams LE. The nonconscious regulation of emotion. In: Gross JJ, editor. *Handbook of Emotion Regulation*. New York: Guilford Press; 2007: 429-455.
3. Campbell-Sills L, Barlow DH. Incorporating Emotion Regulation into Conceptualizations and Treatments of Anxiety and Mood Disorders. In: Gross JJ (Ed) *Handbook of Emotion Regulation*. New York: Guilford; 2007: 542-59.
4. Gross JJ. The emerging field of emotion regulation: An Integrative Review. *Rev General Psychol* 1998; 2(3): 271-99.
5. Eisenberg N, Fabes RA, Guthrie IK, et al. The Relations of Regulation and Emotionality to Problem Behavior in Elementary School Children. *Develop Psychopathol* 1996; 8(1): 141-62.
6. Eisenberg N, Cumberland A, Spinrad TL, et al. The Relations of Regulation and Emotionality to Children's Externalizing and Internalizing Problem Behavior. *Child Development* 2001; 72(4): 1112-34.
7. Mullin BC, Hinshaw SP. Emotion Regulation and Externalizing Disorders in Children and Adolescents. In: Gross JJ, editor. *Handbook of Emotion Regulation*. New York: Guilford Press; 2007: 523-41.
8. Allen LB, Tsao JCI, Seidman LC. A Unified, Transdiagnostic Treatment for Adolescents with Chronic Pain and Comorbid Anxiety and Depression. *Cognitive Behav Pract* 2012; 19(1): 56-67.
9. Aldao A, Nolen-Hoeksema S, Schweizer S. Emotion-Regulation Strategies across Psychopathology: A Meta-Analytic Review. *Clin Psychol Rev* 2010; 30(2): 217-37.
10. Szasz PL, Szentagotai A, Hofmann SG. The Effect of Emotion Regulation Strategies on Anger. *Behavi Res Ther* 2011; 49(2): 114-9.

11. Najmi S, Riemann BC, Wegner DM. Managing Unwanted Intrusive Thoughts in Obsessive-Compulsive Disorder: Relative Effectiveness of Suppression, Focused Distraction, and Acceptance. *Behavi Res Ther* 2009; 47(6): 494-503.
12. Eifert GH, Heffner M. The Effects of Acceptance versus Control Contexts on Avoidance of Panic-Related Symptoms. *J Behavi Ther Experiment Psychiat* 2003; 34(3-4): 293-312.
13. Wetherell JL, Afari N, Rutledge T, et al. A Randomized, Controlled Trial of Acceptance and Commitment Therapy and Cognitive-Behavioral Therapy for Chronic Pain. *Pain* 2011; 152(9): 2098-107.
14. Hayes SC. Acceptance and Commitment Therapy, Relational Frame Theory and the Third Wave of Behavior and Cognitive Therapies. *Behav Ther* 2004; 35(4): 639-65.
15. Lochman JE, Powell NP, Boxmeyer CL, Jimenez-Camargo L. Cognitive-Behavioral Therapy For Externalizing Disorders in Children and Adolescents. *Child Adolescent Psychiatric Clinics North America* 2011; 20(2): 305-18.
16. Bogels S, Hoogstad B, van Dun L, et al. Mindfulness Training for Adolescents with Externalizing Disorders and their Parents. *Behav Cognit Psychotherap* 2008; 36(2): 193-209.
17. Tichovolsky MH. Parenting and Parent Predictors of Changes in Child Behavior Problems. A thesis for Master of Science. University of Massachusetts-Amherst 2011.
18. Wenar C, Kerig P, Ludlow A. *Developmental Psychopathology: From Infancy Through Adolescence*. 6th ed, Maidenhead, UK: McGraw-Hill Education; 2012:
19. Ghomian S, Shairi MR. Preliminary design of the Acceptance and Commitment Therapy Protocol for Children with Chronic Pain (CHACT) and Clinical Trial on the degree of Psychological Inflexibility in Pain of Children 7 to 12 years old. *J Res Behave Sci* 2014; 12(4): 581-92. [in Persian]
20. Minaee A. Adaptation and Standardization Child Behavior Checklist, Self-Evaluation Questionnaire and Teacher's Report Form. *J exceptional Child* 2005; 6(1): 529-58. [in Persian]
21. Potier J, Day C. Childhood onset conduct problems: a preliminary investigation into the role of mothers' interpersonal schemas and their relationship to parenting behaviour. *Behav Cognitive Psychotherapy* 2007; 35(4): 457-72.
22. Strassberg Z. Social Information Processing in Compliance Situations by Mothers of Behavior-Problem Boys. *Child Develop* 1995; 66(2): 376-89.
23. Caspi A, Moffitt TE, Morgan J, et al. Maternal Expressed Emotion Predicts Children's Antisocial Behavior Problems: Using Monozygotic-Twin Differences to Identify Environmental Effects on Behavioral Development. *Developmental Psychology* 2004; 40(2): 149-61.
24. Bronstein P, Duncan P, D'Ari A, et al. Family and Parenting Behaviors Predicting Middle School Adjustment: A Longitudinal Study. *Family Relations* 1996; 45(4): 415-26.
25. Rinaldi CM, Howe N. Mothers' and Fathers' Parenting Styles and Associations with Toddlers' Externalizing, Internalizing and Adaptive Behaviors. *Early Childhood Res Quarter* 2012; 27(2): 266-73.
26. Donnellan MB, Trzesniewski KH, Robins RW, et al. Low Self-Esteem is Related to Aggression, Antisocial Behavior, and Delinquency. *Psychol Sci* 2005; 16(4): 328-35.
27. Espelage DL, Swearer SM. Research on School Bullying and Victimization: What Have we learned and where do we go from here? *School Psychol Rev* 2003; 32(3): 365-83.