Effectiveness of group stress management training on affective control and distress tolerance of mothers of children with sensory-motor disabilities

Original Article

Anahita Khodabakhshi Koolaee (PhD) *1 Soghra Moghimi Nargh (MSc) ²

- Assisstant Professor, Department of Counseling and Psychology, Khatam University, Tehran, IR Iran.
- Msc of Counseling, Rehabilitation Counseling Center, Qum City, Qum, IR Iran.

* Correspondence:

Anahita Khodabakhshi Koolaee, Iran-Khatam University, Hakim Azam St, Shiraz St, Mollasadra Ave. Tehran, 19697-74518, IR Iran.

E-mail:

a.khodabakhshid@khatam.ac.ir

Tel: +98 2189174500 **Fax:** +98 2189174121

Received: 22 Dec 2015 Revised: 19 Jan 2016 Accepted: 10 Feb 2016

Abstract:

Background: The aim of present study was to determine the influence of stress management on affective control and distress tolerance in mothers of children with sensory-motor disabilities.

Methods: This current study was a study with pretest-posttest and treatment group. The thirty mothers of children with sensory-motor disabilities were selected from sensory-motor disabilities center in Qum by 2015. The participants were randomly divided into control (n=15) and experimental (n=15) groups. Stress management training was conducted on experimental group during 8 sessions (90 minutes, twice per week). The research instrument was Affective Control Scale (ACS, 1997) and Distress Tolerance Scale (DTS, 2005). Participants were taken pre-test measures one week prior to the start of this training. Then the questionnaire was administered at post-test. Statistical analysis was performed using analysis of variance.

Results: The results indicated that there was a significant difference between the pretest and post-test scores in distress tolerance (\pm SD: 30.2 \pm 8.1, 33 \pm 9.3; F=7.58) anxiety (\pm SD:40.5 \pm 9.2, 44 \pm 8; F=1.79), anger (SD: \pm 25.4 \pm 4.1, 39.4 \pm 5.6; F=1.03), and depressed mood (SD \pm 27.7 \pm 5.3, 38.6 \pm 7.4; F=1.70) in the experimental group (p<0.05). Additionally, the significant difference was not observed between score of pretest and posttest in positive emotion (SD \pm 41.7 \pm 9.9, 39 \pm 7.48; F=1.08) in the experimental group (p<0.05).

Conclusions: The finding emphasized that the stress management reduced the level of distress tolerance, stress, anxiety, and anger in mothers with sensorymotor disability children. Hence, it can be considered enhancing the affective control in mothers of sensory-motor disability children in therapeutic intervention.

Keywords: Stress/psychological, Affective Symptoms, Adaptation/ psychological, Mothers, Disabled Persons

Citation:

Khodabakhshi Koolaee A, Moghimi Nargh S. Effectiveness of group stress management training on affective control and distress tolerance of mothers of children with sensory-motor disabilities. Caspian J Pediatr March 2016; 2(1): 100-6.

Introduction:

The birth of every child with special needs or behavioral disorders results in several problems for parents. Every child with special needs is born with a unique disability ^[1]. Increasing attention and concentration on disabilities in children lead to personality disorders in parents and children ^[2]. Also, parents have difficulty with accepting the children's disability. Families with disabled children are exposed to greater stresses than other families ^[3]. One of the variables that have strong relation with parent's stress is the severity of disability. Children who are born with severe disabilities cause enormous stress on family ^[4] because family has a primary responsibility for them. There is a distinctive gender role in families with disabled children ^[5].

The presence of mentally retarded child is a source of stress for mothers who involve in caring their children rather than other family member, so it influences their mental health and adaptabilities ^[6]. In other words, mother will be exposed to a danger due to suffering from stress in caring children, relation with others and finally, negative psychological and social effects ^[2]. The disabled child plays a crucial role in the loss of opportunities for social activities and hence it increases the level of stress among these mothers ^[7]. It was distinguished that feeling of tension was related to the decrease of maternal affectionate and responsible behavior; also, the anxiety of mother is associated with disorder in parenting behavior ^[8].

In this regard, Kohsali et al.'s found that there was a significant difference between social adjustments of mothers of mentally retarded daughter and social adjustments of mothers of normal daughters ^[9]. That is why the mean of mental disorder in mothers of exceptional disorder, especially mothers of mentally retarded is higher than normal children's mothers ^[10].

Additionally, families with disabled child have a less health, educational benefits than the others members of society, and this causes that the parents of disabled children report some feelings such as helplessness, low self-esteem and anger [11]. It can be mentioned depression and anxiety as the mental illness and psychological symptoms related to the stress on the mother in this situation. Previous studies indicated that there was a close relation between stress, anxiety and depression [12].

Moreover, anger is a negative predictor of depression in women [13]. In addition, the concept of anxiety is designed by psychologists to explain the responses to failure and dealing with other forms of stress [14]. Therefore, affective control refers to controlling emotions such as stress, anxiety, depression and anger in diverse situations and to ability of accurate expression, so it influence on various aspects of life like interpersonal interactions [15]. Affective control plays an important role in parent-child interaction regarding the negative effect of parent-child interaction in child development [10]. Therefore, stress management can assist disabled child's mother to control stress. Stress management program consists of increasing knowledge in stress, relaxation training, and identification of dysfunctional thoughts, cognitive restructuring, and problem solving and assertiveness skills training [16].

This skill can be reduced the mental pressure, anxiety and daily problems. Emotional expression and

thought process facilitate the emotional adaptability and it leads to changing the attitude toward life, goals and priorities ^[17]. Several researches revealed the efficacy of stress management on reducing stress, anger and anxiety on mental health of the mothers of children with attention deficit hyperactivity disorder ^[18], head-families women ^[19], depressed women ^[20], families with children suffering from hearing-impairment ^[21] and breast cancer women ^[22].

Family training programs provide a context for interaction among parents with common problems. Parents exchange their experiences to help each other so that they can understand children's issues better and attempt to solve them [23] Stress management acquainted individual with stress and coping with it. It can frustrate the effects of stress and stress response, and finally it helps individual to have a better physiological and psychological function. This intervention can be effective through increasing the sense of control, self-efficacy, self-esteem, adaptive coping and social support [24]. Farahani et la. showed the efficacy of stress management group training on aggressiveness [25]. Shokohi and Zamani indicated that anger management training led to increasing the positive relation of mothers with mentally retarded children and slow learners [26].

However, the influence of stress management training on tolerance and affective symptom on mothers with sensory-motor disability children were not studied in previous researches such as Farahani et al.'s [25] and Shokohi et al.'s [26].

Therefore, this study was conducted for the following reasons; firstly, since any disability in affective control and management affects directly on the mental and physical health of mothers. In addition, the efficacy of stress management on reducing of psychological problems has shown in other samples. Finally, women have considerable influence on family and especially on disabled child. Thus, this research intended to respond to this question: what is the influence of stress management training on affective control and distress tolerance of mothers of children with sensory-motor disabilities?

Methods:

This study was conducted in 2015 in Qum. The present research is a study with pre-posttest and treatment group. The sample group was 30 mothers of disabled children, who interested to the training program. Mothers were selected through convenience

sampling method according to inclusion criteria from sensory-motor disabilities center in Qum by 2015. These samples were randomly divided into treatment and control groups. Stress management (table 1) was implanted in 8 seasons (90 minutes for each season, twice a week) for the treatment group based on the theory of cognitive - behavioral stress management provided by Anthony et al.'s [24], but the control group did not receive any intervention.

The researcher followed the standardized procedures and techniques of the training program accurately. To minimize the environmental difference between research and control groups, the same-trained psychologist instructed the groups and the sessions were held in the same place. Part of each session was devoted to review the content, topics and assignments, from previous session. At the end of each session, the assignment was given for next therapy session.

After the intervention sessions, the post-test was taken for experimental and control groups. The inclusion criteria were as follow: mothers had to have a child with sensory-motor disabilities, high school diploma, mothers did not have any severe mental and physical illnesses. Exclusion criteria were absence more than one session and participation in other psychological interventions.

To respect the rights of participants, some information was provided about the performance of program and the researcher emphasized on respect the principal of the confidentiality of personally information and written forms.

Informed consent was obtained from parental or guardian before the actual training started. Williams et al.'s developed the Affective Control Scale (ACS) [27] to assess fear of losing control over one's emotions or of one's behavioral reactions to emotion.

The 42-items are rated on 7-point Likert-type scales from 1 (very strongly disagree) to 7 (very strongly agree) and compose four subscales: fear of anger, depression, anxiety, and positive emotion. The total score is the average of all items. This scale was applied for 105 introductory psychology students from American University.

Internal consistency was satisfactory for the total score (Cronbach's alpha=0.94) as well as for the subscale scores (0.72 anger, 0.91 depression, 0.89 anxiety, 0.84 positive affect). Test-retest reliability for the total score was acceptable (r=0.78). Tahmasebian et al.'s. [15] normalized this scale. They carried out the test on all students, teachers, nurses (n=1500) in Kermanshah. In Persian study, the alpha cronbach

reliability in students, teachers, and nurses was 0.78, 0.88, and 0.90 respectively.

Distress tolerance scale was designed by Simons and Gaher ^[28] to assess the extent to which individuals experience negative emotions without acting to avoid, alleviate, or become absorbed in them. This inventory is 15-item-self-report measures that reflect perceived ability to tolerate emotional distress, subjective appraisal of distress, attention being absorbed by negative, and regulation efforts to alleviate distress. Items were rated on a 5-point scale from ^[5] strongly disagree to ^[1] strongly agree. High scores represent high distress tolerance.

It was conducted on 642 students recruited from two state universities. Internal consistency was satisfactory for the total score (Cronbach's alpha=0.81) as well as for the subscale scores (0.71 tolerance, 0.69 absorption, 0.77 appraisal, 0.73 regulation).

Test-retest reliability over a 6-month interval was good (r=0.61). Moreover, in Persian study, it was performed among mothers of premature infants in Mashhad in 2008 and the Cronbach's alpha of total score was 0.82 and the subscale scores were 0.72, 0.82, 0.78, 0.70 tolerance, absorption, appraisal and regulation, respectively [8].

The collected data were analyzed using SPSS-16. Finally, data were compared between these two groups using ANCOVA.

Table 1. Structure of stress management skills

Sessions	Sessions Objectives						
First	Initial introduction, declare short						
Second	Relaxation training						
Third	Automatics thoughts and Cognitive distortions						
Forth	Self-Talk and thought control strategies						
Fifth	Concentration and thoughts, Distraction						
Sixth	Problem solving training						
Seventh	Anger management training						
Eighth	Increasing self-esteem to adapt with their current situation and training self- assertiveness						

Results:

Table 2 indicated the results of socio-demographic of all mothers participated in this study. Totally, ten (66.68%) and eight (53.33%) mothers had diploma in experimental and control groups, respectively. The most numbers of mothers in experimental (33.3%) and control (46.66%) groups were the same age (36-40 years). Mean scores of pre-test and post-test in two

groups are shown in table 3.In addition, it was utilized the Leven test to verify equal variances (homogeneity of variance) and normally distributed.

Table 4 shows the results of variance analysis to compare the meaning of these two groups by eliminating the effectiveness of pre-test.

According to these findings, there was a significant difference between the scores of pre-test and post-test

Table 2. General characteristic of mothers' disabled children

Variable	Group				
variable	Experimental	Control			
Age (y)	N (%)	N (%)			
25-30	1 (6.66)	2 (13.33)			
31-35	4 (26.6)	3 (20)			
36-40	5 (33.3)	7 (46.66)			
41-45	5 (33.3)	3 (20)			
Education					
High School	2 (13.33)	3 (20)			
Diploma	10 (66.68)	8 (53.33)			
Baccalaureate Degree	2 (13.33)	1 (6.66)			
Master's	1 (6.66)	3 (20)			

in distress tolerance (F=7.58, p=0.014, η 2=433), anxiety (F=1.79, p=0.001, η 2=0.90), anger (F=1.03, p=0.002, η 2=0.92), and depressed mood (F=1.70, p=0.011, η 2=0.91). Moreover, the significant difference was not observed between the scores of pretest and post-test in positive emotion (F=1.08, p=0.17, η 2=0.91).

Table 3. Mean scores of pre-test and post-test in two groups of participants

Variable	Groups	Pre-test M (SD)	Post-test M (SD)
Distress	Experi	33.60 (8.5)	30.2 (8.1)
tolerance	Control	35 (11.43)	33 (9.3)
Anxiety	Experi	44.20 (3.4)	40.5 (9.2)
	Control	46.1 (7.64)	44 (8)
Anger	Experi	26.10 (3.3)	25.4 (4.1)
	Control	27.2 (5.2)	39.4 (5.6)
Depressed	Experi	37.5 (3.4)	27.7 (5.3)
mood	Control	39.4 (4.33)	38.6 (7.4)
Positive	Experi	37.6 (8.41)	41.7 (9.9)
emotion	Control	39.3 (7.18)	39 (7.48)

N= 30 (Experimental group=15, Control Group= 15)

Table 4. The comparison of the total dimension of quality of life in patients' age, education and disease stage

variable	Group	^b SS	^b Df	^b MS	F	P	b η^2
Distress tolerance	Covariate	137.33	1	137.33	7.58	0.014	0.70
	Between	559.63	1	559.64			
	Residual	1180.42	13	73.77			
Anxiety	Covariate	755.72	1	755.72	1.79	0.001	0.90
	Between	107.74	1	107.74			
	Residual	962.71	13	60.17			
Anger	Covariate	15.97	1	15.97	1.03	0.002	0.92
	Between	111.68	1	111.68			
	Residual	26.71	13	15.42			
Depressed mood	Covariate	47.84	1	47.84	1.70	0.011	0.91
	Between	58.47	1	58.47			
	Residual	549.84	13	34.36			
Positive emotion	Covariate	898.4	1	898.4	1.08 0.17		0.91
	Between	2.34	1	2.34		0.17	
	Residual	458.29	13	28.46			

^bAbbreviations: SD, Standard Deviation; df, degree of freedom; SS, Sum of Square; MS, Means of Square; η2, Eta Square

*P<0.05

Discussion:

The current study was conducted to investigate the influence of stress management training on the distress tolerance and emotional control among mothers of sensory-motor disabled children. Based on this finding, stress management skills elevate the distress tolerance among mothers of sensory-motor disabled children. This result is consistent with some prior researchs [29, 30, 31]

Harris et al.'s was investigated the influence of stress management (yoga-based) on sixty-four educators in two middle schools. Intervention participants had significantly improved scores on distress tolerance ^[29]. Gawrysiak et al.'s showed that Mindfulness-Based Stress Reduction decline in perceived stress through influence on distress tolerance. Additionally, individuals with lower baseline distress tolerance evidenced a greater decline in perceived stress ^[30].

In addition, Alavi et al.'s showed that group dialectical behavior therapy (based on core mindfulness and emotion regulation components) reduced the distress tolerance among university students [31]. Distress tolerance is conceptualized as a metaregulation process of how individuals react to uncomfortable emotions. Therefore, stress management help individual to recognize the situation which causes the anxiety and stress, then they learn coping strategies to coping with stress. This process leads to reduce distress tolerance of mothers of sensory-motor disabilities children who encounter obstacle during early child development in diverse aspects.

Additionally, the findings of current study revealed that stress management increased the level of affective control. Therefore, it reduced the level of anxiety, anger and depressed mood in mothers of sensory-motor disabilities children. These results were consistent with prior study in this field [32], For example, Akhteh et al.'s showed that stress management reduced the level of anxiety in women with recurrent miscarriage [33]. Marsland et al.'s. revealed that stress management influenced on acceptability and feasibility of mothers of children newly diagnosed with cancer. It could reduce the level of distress in this group [34]. Findings of Parand and Mollavi showed a significant reduction of psychological problems (stress, anxiety, and anger) in mothers after participating in the stress management program [21]. Dehghan et al.'s investigated the influence of stress management in reducing of aggressiveness among students. They found that there were significant

differences in per- and post- test scores for all aggression components in experimental group [35].

The result of current study suggested that there was no relationship between positive affect and stress management training. The positive affect is referred to feeling happy, proud, and the level of tendency for life [36]. The previous study revealed that this item was provided by social support of family and friends among adult with disabilities [37] or by increasing the level of meaning of life through hope therapy among mothers of sensory-motor disabled children [38]. Consequently, it can be concluded that stress management does not play role in increasing the level of the positive affect.

It should be considered that one of the factors which influence on elevating the level of stress and anxiety is negative thoughts. Heevady et al.'s. found that irrational believe of blame proneness, frustration reactive, anxious over concern, problem avoiding, dependency and perfectionism among mothers of sever or profound mentally handicap child were more than among mothers with normal child [39]. The relation with irrational belief and anger, anxiety and stress was approved in previous studies [40]. Thus, stress management program relieves anger, depressed mood and anxiety through the reduction of negative thoughts and self-talk. In other words, since the parents of children with sensory-motor disabilities increasingly at risk of stress and other problems related to mental health, the growth of self-awareness (through the reduction of negative thoughts) and coping strategies should be considered important to increase the distress tolerance and affective control on parents.

Like our study, Khodabakhshi Koolaee et al.'s. also indicated there was statistically significant difference between mother with or without children with speech disorder [41] and with mental retardation [42] in caregiver burden.

Limitation of this study was the use of self-report measure which increased the possibility of biased reports. In addition, the generalizability of this finding should be done cautiously due to the limited sample (mothers with disability children) in Qum. Further studies should include more than one family member.

Recommendation:

Therefore, the results of current and previous research emphasize that the intervention program aimed to reduce the stress of parents who have children with mental disabilities is necessary.

Acknowledgment:

Thanks to all mothers of disability children, who participated in this study.

Funding: This study was self-funded.

Conflict of interest: There was no conflict of interest.

References:

- Khodabakhshi koolaee A. family therapy and parent training; program & models: A comprehensive guide to solve problems behaviors in children and adolescents. 2th ed. Tehran: Jangal Publication. 2012: 8-10. [in Persian]
- Ahmadi Kh A, Khodadadi Gh, Anisi J, Abdolmohammadi E. Problems of families with disabled children. J Mil Med 2011; 13(1): 49-52. [in Persian]
- 3. Woodman AC. Trajectories of Stress among Parents of Children with Disabilities: A Dyadic Analysis. Family Relations 2014; 63(1): 39-54.
- Cohen E, Yantzi N, Guan J, et al. Residential movement patterns of families of young children with chronic conditions in Ontario, Canada: a population-based cohort study. Int J Equity Health. 2013; 12: 62.
- 5. Lindsay S, King G, Klassen AF, et al. Working with immigrant families raising a child with a disability: challenges and recommendations for healthcare and community service providers. Disabil Rehabil 2012; 34(23): 2007-17.
- Sulkers E, Tissing WJ, Brinksma A, et al. Providing care to a child with cancer: a longitudinal study on the course, predictors, and impact of caregiving stress during the first year after diagnosis. Psychooncology 2015; 24(3): 318-24.
- Benson PR. Longitudinal effects of educational involvement on parent and family functioning among mothers of children with ASD. Research in Autism Spectrum Disorders 2015; 11: 42-55.
- 8. Reihani T, Sekhavat Poor Z, Heidarzadeh M, et al. The Effect of Spiritual Self-Care Training on Feeling of Comfort in Mothers of Hospitalized Preterm Infants. J Midwifery Reproductive Health 2014; 2(2): 112-9. [in Persian]
- 9. Koohsali M, Mirzamani SM, Karimlo M, Mirzamani MS. Comparison of social adjustment in mothers of educable mentally retarded daughter. J Behav Sci 2008; 2(2): 165–171. [in Persian]
- 10. ZandvanianNaeini A, Morowati sharifabad MA, Amrollahi M. Efficacy of Anger Management Training on increasing Mental Health of Disabled Children Mothers. TB 2014; 13(3): 73-85.

- 11. Ahmadi K, Khodadadi G, Akhavi Z, Anisi J. Function of families with disabled children. J Behav Sci 2013; 6(4): 331-7. [in Persian]
- 12. Uliaszek AA, Zinbarg RE, Mineka S, et al. A longitudinal examination of stress generation in depressive and anxiety disorders. J Abnorm Psychol 2012; 121(1): 4-15.
- 13. Rude SS, Chrisman JG, Burton Denmark A, Maestas KL. Expression of direct anger and hostility predict depression symptoms in formerly depressed women. Canadian J Behav Sci 2012; 44(3): 200-9.
- 14. Servat F, Morowatisharifabad M, Sadeghipoor M, et al. Study of the Effect of Stress Management Skills on Anxiety Status among Teachers in Meybod District High Schools On 2012. J Toloo e Behdasht 2015; 14 (1): 89-100. [in Persian]
- 15. Tahmasebian H, Khazaie H, Arefi M, et al. Normalization of emotion control scale. J Kermanshah Uni Med Sci 2014; 18(6): 349-54. [in Persian]
- Lehrer PM, Woolfolk RL, Sime WE. Principles and practice of stress management. Guilford Press; 3th ed. NY: Guilford Press; 2007: 38-50.
- 17. Jamilian H, Khansari M, Safari V. The Comparison of Efficacy of "Stress Management Training Through Cognitive Behavioral Method" and "Relationship Skills Training with End-Of-Life Patient" on Increase of Family Life Quality. J Urmia Uni Med Sci 2014; 24(12): 987-95. [in Persian]
- 18. Zarei Sh, Sharif F, Alavi Shooshtari A, Vossoughi M. The effect of stress management program on mental health of the mothers of children with attention deficit hyperactivity disorder. J Nurs Edu 2015; 3(4):58-67. [in Persian]
- 19. Habibi M, Ghanbari N, Khodaei E, Ghanbari P. Effectiveness of cognitive-behavioral management of stress on reducing anxiety, stress, and depression in head-families women. J Res Behav Sci 2013; 11(3): 166-75. [in Persian]
- 20. Abbasian F, Najimi A, Ghasemi Gh, et al. Effectiveness of Stress Management Training by Cognitive-Behavioral Method in Women with Depression. J Health System Res 2013; 8(6): 1050-57. [in Persian]
- 21. Parand A, Movallali G. The effect of teaching stress management on the reduction of psychological problems of families with children suffering from hearing-impairment. J Family Res 2011; 7(1): 23-34. [in Persian]
- 22. Khodabakhshi Koolaee A, Falsafinejad MR, Esmaeili Akbari M. The Effect of Stress

- Management Model in Quality of Life in Breast Cancer Women. Iran J Cancer prev 2015; 8(4): e3435.
- 23. Parand A, Afrooz Gh, Mansoor M, et al. Developing stress management program for mothers of children with ADHD and its effectiveness on their mental health. Procedia-Soc Behav Sci 2010; 5: 1135-9.
- 24. Antony M, Ironson G, Eshnaiderman N. A Practical Guide stress management program utilizing cognitive-behavioural approach. Translator: Alemohammad J, Jukar S, Neshatdost H. 1th ed: Esfahan: Publications Academic Jihad; 2009: 48-64. [in Persian]
- 25. Farahani M, Basaknejad S, Davoodi I. Effectiveness of stress control training group on aggression, parental stress and mental health of mothers of children with learning disabilities. Psychological Achievements 2013; 20(2): 217-218 [in Persian].
- 26. Shokohi Yekta M, Zamani N. Efficacy of Anger Management Training on Relationship Between Mothers and Their Mildly Mentally Retarded and Slow Learner Children. J Family Res 2008; 4(15): 231-46. [in Persian]
- 27. Williams KE, Chambless DL, Ahrens A. Are emotions frightening? An extension of the fear of fear construct. Behav Res Ther 1997; 35(3): 239-48.
- 28. Simons JS, Gaher RM. The Distress Tolerance Scale: Development and validation of a self-report measure. Motivat Emot 2005; 29(2): 83-102.
- 29. Harris AR, Jennings PA, Katz DA, et al. Promoting Stress Management and Wellbeing in Educators: Feasibility and Efficacy of a School-Based Yoga and Mindfulness Intervention. Mindfulness 2016; 7(1): 143-54.
- 30. Gawrysiak MJ, Leong SH, Grassetti SN, et al. Dimensions of distress tolerance and the moderating effects on mindfulness-based stress reduction. Anxiety Stress Coping 2015: 1-9.
- 31. Alavi Kh, Modaress Gharavi M, Amin Yazdi SA, Salehi Fadardi J. Effectiveness of group dialectical behavior therapy (based on core mindfulness, distress tolerance and emotion regulation components) on depressive symptoms in university students. J Fundamental Mental Health 2011; 13(2): 124-35. [in Persian]
- 32. Brown FL, Whittingham K, McKinlay L, et al. Efficacy of Stepping Stones Triple P plus a stress management adjunct for parents of children with an acquired brain injury: The protocol of a randomised

- controlled trial. Brain Impairment 2013; 14(2): 253-69
- 33. Akhteh M, Alipor A, Sharifi Saki Sh. The effectiveness of stress management on anxiety and meta-worries women with recurrent miscarriage. Health Psychology 2014; 3(11): 119-29. [in Persian]
- 34. Marsland AL, Long KA, Howe C, et al. A pilot trial of a stress management intervention for primary caregivers of children newly diagnosed with cancer: preliminary evidence that perceived social support moderates the psychosocial benefit of intervention. J pediatr psychol 2013; 38(4): 449-61.
- 35. Dehghan F, Karami J, Piri M, Karimi P. The Efficiency of Stress Management Skills Training in Reducing Aggression. J Mazandaran Univ Med Sci 2014; 24(115): 162-166 [in Persian]
- 36. Pintrich PR. Multiple goals, multiple pathways: The role of goal orientation in learning and achievement. J Edu Psychol 2000; 92(3): 544-55.
- 37. Imani M, Khodabakhshi Koolaee A, Rahmatizadeh M. Association between social support of family and friends and meaning of life with Depression among spinal cord injuries disabilities and non-disabilities. Med East J Psychiatr Alzheimers 2013; 4(3): 8-12.
- 38. khodabakhshi koolaee A, Derakhshandeh M. Effectiveness of hope-oriented group therapy on life meaning and resilience in mothers with physical-motor disabled children. J Pediatr Neonatal Nurs 2015; 1(3):15-25. [in Persian]
- 39. Hivadi B, Mirzayi SM, Bahrami H, Salehi M. Comparison of Irrational Believes between Mothers of Severe or Profound Mentally Handicapped Children with Healthy Children Mothers. J Rehabil 2007; 8(3): 75-80. [in Persian]
- 40. Fives CJ, Kong G, Fuller JR, DiGiuseppe R. Anger, Aggression, and Irrational Beliefs in Adolescents. Cognitive therapy and research 2011; 35(3): 199-208.
- 41. Khodabakhshi Koolaee A, Baghersad H, Rahmatizadeh M. The psychological effects of having a child with a speech disorder. Middle East J Psychiatry Alzheimers 2014; 5(1): 3-8. Available at: http://platform.almanhal.com/Article/Preview.aspx?I D=13718
- 42. Khodabakhshi Koolaee A, Khazan S, Tagvaee D. Mother-Child relationship and burden in families of children with mental retardation. Middle East J Psychiatry Alzheimers 2014; 5(3): 32-6.