

# Glucose-6-phosphate dehydrogenase enzyme deficiency in Iranian newborns: A systematic review and meta-analysis

## Review Article

Yadollah Zahed Pasha (MD)<sup>1</sup>

Vahid Zamani (MD)<sup>2</sup>

Ermia Zahed Pasha (MD)<sup>3</sup>

Mohammad Zamani<sup>4, 5 \*</sup>

1. Non-Communicable Pediatric Diseases

Research Center, Health Research Institute,  
Babol University of Medical Sciences, Babol,  
Iran.

**ORCID ID** orcid.org/0000-0002-3011-0361

2. Vice-Chancellery for Health, Babol

University of Medical Sciences, Babol, Iran.

3. Department of Pathology, Rouhani Hospital,

Babol University of Medical Sciences, Babol,  
Iran.

4. Student Research Committee, Babol

University of Medical Sciences, Babol, Iran.

**ORCID ID** orcid.org/0000-0003-1916-3873

5. Cancer Research Center, Health Research

Institute, Babol University of Medical  
Sciences, Babol, Iran.

### \* Correspondence:

**Mahammad Zamani**, Ganjafrouz Street,  
Babol University of Medical Sciences, Babol,  
47176-47745, I.R. Iran.

**E-mail:** mzamani20@gmail.com

**Tel:** +98 1132190971

**Fax:** +98 1132190971

**Received:** 12 Dec 2019

**Revised:** 25 Jan 2020

**Accepted:** 18 Feb 2020

## Abstract

**Background:** The aim of this study was to perform a systematic review and meta-analysis on available data about glucose-6-phosphate dehydrogenase enzyme deficiency (G6PDD) status in Iranian neonates screened for the disease.

**Methods:** A literature search was conducted in electronic databases of Embase, PubMed, Web of Sciences, Scopus and Google Scholar for articles published from inception to 1 December 2018. Moreover, the literatures from Iranian databases, including Magiran and Scientific Information Database were searched. We included observational studies reporting prevalence of G6PDD, related complications and genetic factors among Iranian neonates. Data were analyzed using STATA software.

**Results:** Of 656 articles were initially found, 16 were finally included. Overall pooled prevalence of G6PDD was 5.5% (95% confidence interval: 2-8.9). Analysis also indicated that boys were significantly 3 times more at risk of G6PDD compared with girls. Three articles were identified related to the jaundice and 4 papers related to kernicterus. A range of 43-67% of newborns with G6PDD presents with jaundice. Additionally, 5-9% of G6PDD cases with jaundice present with kernicterus. One article reported that out of 412 newborns, 12.9% were carriers for one of the three G6PD gene mutations, including Mediterranean, Chatham and Cosenza.

**Conclusions:** Prevalence of G6PDD in Iran is comparable to most countries. Jaundice and kernicterus are major complications of G6PDD. Therefore, it is necessary to pay attention to all patients with G6PDD. Also, it is recommended that hospitals provide the result of G6PD testing as soon as possible and before discharging newborn children.

**Keywords:** Glucose-6-phosphate dehydrogenase enzyme deficiency, G6PD, Hemolytic anemia, Jaundice, Kernicterus

## Citation:

Zahed pasha Y, Zamani V, Zahed pasha E, Zamani M. Glucose-6-phosphate dehydrogenase enzyme deficiency in Iranian newborns: A systematic review and meta-analysis. *Caspian J Pediatr* March 2020; 6(1): 376-86.

## Introduction

Glucose-6-phosphate dehydrogenase enzyme deficiency (G6PDD) is an X-linked genetic disease caused by mutations in the G6PD gene, and is the most common enzymopathy in the world [1]. It is reported that this disease worldwide affects approximately 400 million people and 11 million infants with G6PDD, born every year [2, 3]. G6PDD has a prevalence of 5-25% in areas where malaria is endemic, and <5% in non-endemic areas [4]. G6PD enzyme exists in all body cells and has an important role in protecting against oxidative stress. In the patients with G6PDD, due to its oxidation of the red blood cell membrane, the cells are destroyed, causing hemolysis [5].

Although G6PDD is usually asymptomatic, it can have serious clinical features –that is, hyperbilirubinemia and jaundice, both of which result from an increased rate of hemolysis [6, 7]. One of the important risks of hyperbilirubinemia is kernicterus, which can cause irreversible neurologic complications and permanent developmental disorders observed more frequently in neonates [8, 9]. Because the complications of G6PDD are more serious in newborns, it is necessary to determine the presence of this genetic disease early in this age group.

Health care systems around the world consider a screening program as an important and efficient step in the reduction of hospitalization caused by favism (a form of hemolytic anemia caused by contact with broad beans) and also kernicterus. Most of countries have this program in neonates [10, 11]. In Iran, there is a newborn screening program for G6PDD and it is managed by the Ministry of Health and Medical Education [12]. We aimed to perform a systematic review and meta-analysis on the available data about G6PDD prevalence in Iranian neonates screened for the disease. In addition, it was tried to collect the information related to the genetic factors associated with G6PDD and complications which occur following the disease. These data should be useful for clinicians and other health professionals planning for better management of G6PDD in Iranian newborns.

## Methods

**Information sources and search strategy:** A literature search was conducted in the electronic databases of Embase, PubMed, Web of Sciences, Scopus and Google Scholar for articles published from inception to 1 December 2018. After searching the related terms in the Medical Subject Headings (MeSH) database, finally, the keywords included “glucose-6-phosphate dehydrogenase deficiency” OR “Glucosephosphate Dehydrogenase Deficiency” OR “Glucose 6 Phosphate Dehydrogenase Deficiency” OR “G6PD deficiency” AND “Iran” OR “Iranian”. The search was limited to Title/Abstract. The word “Iran” was limited to Affiliation as well. Furthermore, the current study searched literatures from Iranian databases, including Magiran and Scientific Information Database (SID), using the Persian equivalent of the above-mentioned keywords. Hand searching was also performed on the reference lists of the relevant review articles and studies finally included in the current study to identify additional sources.

This systematic review and meta-analysis were conducted according to the guideline of Preferred reporting items for systematic review and meta-analysis (PRISMA) [13]. The protocol of the present study is available in the PROSPERO registry, too (CRD42019119693) [14].

**Inclusion and exclusion criteria:** We included observational studies reporting the prevalence of G6PDD among Iranian neonates. To have an acceptable and real prevalence, we included the studies screening for G6PDD in newborns, but we excluded from further analyses conducted only on the subjects with jaundice/hyperbilirubinemia or any other specific disease. We included the later studies for assessment of complications of G6PDD and genetic factors potentially associated with the disease. The other exclusion criteria were the following:

1. Reviews, case reports, editorials, letters and comments,
2. Duplicate articles,
3. Studies conducted on subjects other than neonates,
4. Articles without clear methodology or results, and Full-texts were not available.

**Study selection and data extraction:** Two authors (MZ, VZ) assessed the Titles and Abstracts independently for eligibility. Besides, the full-text of the potential articles was evaluated in the next step. When there was a discrepancy, it was resolved by consensus with a third author (YZ). Two authors (MZ, EZ) extracted data independently. The following data were collected: first author's name, study period, publication date, study location, number of population (total and by gender), prevalence of G6PDD among screened neonates, prevalence of complications in newborns with G6PDD. Duplicate articles were excluded and one with more details or larger sample size was selected.


**Quality assessment:** The checklist by Hoy et al. [15] was used for evaluating the risk of bias, which has nine questions with two potential responses (Yes/No). The range of scores is between 0 and 9. Higher scores are representative of higher risk of bias.

**Study outcomes and statistical analysis:** After collecting the necessary data, they were analyzed using STATA software (StataCorp, College Station, TX, USA). The pooled estimate rate of G6PDD prevalence was presented as percent and 95% confidence interval (CI). The complications of G6PDD in the present study were jaundice (in neonates with G6PDD) and kernicterus (in G6PDD cases presented

with jaundice). Additionally, the sub-group analyses were performed by gender (male and female) and study date (<2007 and ≥2007). Splitting the study date into <2007 and ≥2007 was mainly based on the distribution of the number of reports in each period category. When the study date was not mentioned, the year of study publication was used instead. Prevalence of G6PDD was compared according to gender by using an odds ratio (OR) with a 95% confidence interval (CI). I<sup>2</sup> statistic and  $\chi^2$  test were used for checking the statistical heterogeneity. Random effects model was used for meta-analysis. Forest plots were provided to summarize the results of meta-analyses.

## Results

A total of 656 articles were initially found by searching the databases, of which 198 papers were excluded after evaluating title/abstract. After assessing full-texts of 35 articles, 19 studies were excluded. Different steps of systematic review were indicated in PRISMA chart (Figure 1). Overall, 16 studies were included in the systematic review for final analysis and their details were summarized in table 1.



**Figure 1. Prisma Flowchart**

**Table 1. Characteristics of the studies reporting prevalence of G6PDD among Iranian newborns screened for the disease**

Region	Author	Publication date	Study date	Risk of bias score	Number (total)	Number (boys)	Number (girls)	Prevalence (%; total)	Prevalence (%; boys)	Prevalence (%; girls)
<b>Babol</b>	Zahedpasha (54)	1999	1995	1/9	2046	1035	1011	8.3	12.5	4.1
<b>Bushehr</b>	Movahhed (55)	2003	1998	1/9	415	218	210	8.4	12.8	1.9
<b>Fars</b>	Daliri (56)	2017	2011-2015	1/9	383463	199536	183927	15.6	16.3	14.9
<b>Isfahahn</b>	Iranpour (57)	2008	2006	1/9	2501	1307	1194	3.2	5.1	1
<b>Mashhad</b>	Mohammadzadeh (58)	2009	2006	1/9	2570	1307	1263	0.8	1	0.5
<b>Mazandaran</b>	Kosaryan (59)	2011	2007-2010	1/9	115622	59430	56192	5.8	-	-
<b>Rafsanjan</b>	Alidalaki (60)	2007	2004	1/9	1018	523	495	5	5.7	4.2
<b>Rasht</b>	Khalili (61)	2007	2001	1/9	1197	605	585	6.4	9.8	3.1
<b>Rasht</b>	Keihanian (62)	2016	-	1/9	1474	757	717	6.6	11.4	1.7
<b>Sari</b>	Kosaryan (63)	2014	2012-2013	1/9	365	174	191	7.5	7.5	0
<b>Semnan</b>	Nazari (64)	2011	2007-2010	1/9	9353	4820	4533	3.2	5.5	0.8
<b>Shahrekord</b>	Norbakhsh (65)	2013	2011	1/9	1240	633	607	2.3	2	2.6
<b>Sistan and Baluchestan</b>	Ansari-Moghaddam (66)	2017	-	1/9	140	68	72	8.4	11.8	5.6
<b>Tehran</b>	Abolghasemi (17)	2004	1999	1/9	2000	1006	994	2.1	3.6	0.6
<b>Tehran</b>	Khalesy (16)	2012	2008-2009	1/9	450	245	205	2	3.3	0.5
<b>Tehran</b>	Kazemi (18)	2013	2009	1/9	1226	585	641	2.2	2.1	2.3


**Table 2. Rate of jaundice in neonates with glucose 6-phosphate dehydrogenase deficiency and kernicterus in those presented with jaundice in Iran**

Complication	Author	Number of subjects	Prevalence of complication (%)
Jaundice	Abolghasemi (17)	42	42.9
	Kazemi (18)	27	70.3
	Khalesy (16)	9	66.7
Kernicterus	Aletayeb (19)	112	4.5
	Boskabadi (20)	59	6.8
	Boskabadi (22)	45	8.9
	Yousefi (21)	34	8.8


### Prevalence:

Analysis of 16 articles showed that the overall prevalence of G6PDD in Iran was 5.5% (95% CI: 2-8.9) (Figure 2). The highest rate was for Fars (15.6%) and the lowest one pertained to Mashhad, Razavi Khorasan (0.8%). In addition, the overall pooled prevalence of G6PDD was 7.3% (95% CI: 2.8-11.8) in boys and 3.1% (95% CI: 0-7.6) in girls (Figure 3). Subgroup analysis by study date indicated that the


overall estimated prevalence of G6PDD was 4.8% (95% CI: 2.8-6.7) based on studies conducted before 2007, and 5.9% (95% CI: 1.4-10.5) based on studies performed after 2007 (Figure 4). Analysis also suggested that boys were significantly 3 times more at risk of G6PDD than girls (OR=3.1, 95% CI: 1.8-5.3) (Figure 5).



**Figure 2. The overall pooled prevalence of G6PDD in Iranian newborns**




**Figure 3. The overall pooled prevalence of G6PDD in Iranian newborns by gender**



**Figure 4. The overall pooled prevalence of G6PDD in Iranian newborns by study date**





**Figure 5. Pooled Odds Ratio (OR) for G6PDD in boys compared with girls**

### Complications:

In the present systematic review, also it was tried to find the data on the complications of G6PDD among Iranian newborns, including jaundice and kernicterus. After searching the databases, 3 articles [16-18] were identified related to the jaundice and 4 papers [19-22] related to the kernicterus (Table 2). Based on the reports, it was determined that a range of 43-67% of newborns with G6PDD presented with jaundice. In addition, 5-9% of G6PDD cases with jaundice presented with kernicterus, too.

### Genetic factors:

Limited number of articles was found that reported genetic factors in G6PDD Iranian newborns. There was only one article assessing the frequency of G6PD mutations, which was carried out in the North of Iran. Mahdavi et al [23]. Stated that 12.9% of 412 newborns (8.6% of boys and 16.8% of girls) were carriers for one of the three G6PD gene mutations including Mediterranean, Chatham and Cosenza. The Mediterranean type was the most frequent mutation among the three examined molecular mutations.

One study by Zahedpasha et al [24]. In Northern Iran investigated the association between the three mutations of G6PD and jaundice. After comparing the

distribution of Mediterranean and Chatham mutations between icteric and non-icteric neonates (both with G6PDD), no any significant differences were recognized. On the other hand, the distribution of rare mutations (Cosenza negative) was significantly higher in non-icteric than in icteric newborns. Some mutations of G6PD gene may less likely lead to neonatal icterus, for example, neonates with the rare Chatham mutation are less in need of exchange transfusion,

A different survey by Zahedpasha et al [25]. evaluated any possible relation between neonatal icterus and Gilbert syndrome in newborns with G6PDD, but no any significant differences were found between icteric and non-icteric subjects in the distribution of Gilbert syndrome.

### Discussion:

In this systematic review and meta-analysis, it was determined that the prevalence of G6PDD among Iranian newborns was 5.5%. We included the studies in which the newborns underwent screening for G6PDD. There are different studies about G6PDD prevalence in neonates worldwide. For instance, the screening studies on neonates demonstrated G6PDD rates as 11.1% in

the United States, <sup>[26]</sup> 4.5% in India, <sup>[27]</sup> 6.1% in Thailand, <sup>[28]</sup> 2.4% in China, <sup>[29]</sup> 7.8% in Brazil, <sup>[30]</sup> 4.3% in Egypt, <sup>[31]</sup> 4.4% in Tunisia <sup>[32]</sup> and 2% in Saudi Arabia <sup>[33]</sup>. Variable prevalence between different countries and regions can be explained by differences in ethnicity and genetic distribution between populations. Many G6PD gene mutations are responsible for deficiency of the enzyme, including Mediterranean, Chatham, Cosenza, and G6PD A and so on. In a systematic review, only one article was found that reported on the prevalence of G6PD mutations among Iranian newborns <sup>[23]</sup>.

In a recent meta-analysis concerning Iran, the prevalence of Mediterranean, Chatham, and Cosenza molecular mutations was estimated 78.2, 9.1 and 0.5% in G6PD-deficient people, respectively <sup>[34]</sup>. Mediterranean G6PD has a high prevalence in other tropical and subtropical regions <sup>[24]</sup>. Its prevalence is similarly high in neighboring countries, such as Saudi Arabia (80%), Oman (74%), Turkey (77%), India (60.4%), United Arab Emirates (55.5%) and Pakistan (76%), as well as in Mediterranean coast countries <sup>[34]</sup>. This mutation is mainly associated with favism <sup>[35, 36]</sup>. Chatham mutation is responsible for G6PDD class II presenting with severe hemolytic anemia; however, it has lower prevalence compared with the Mediterranean mutation <sup>[37-39]</sup>.

Our subgroup analysis showed a 3-fold higher rate of G6PDD in boys than in girls and risk of G6PDD, a result that was consistent with previous research <sup>[40]</sup>. Considering that inheritance G6PDD has an X-linked pattern, it is expected to see this disease more in male infants than in female ones. Homozygous males with class I mutations usually develop chronic non-spherocytic hemolytic anemia, whereas females who are heterozygous for G6PD can carry severe mutations but may remain symptomless <sup>[41, 42]</sup>.

Neonatal screening for G6PDD is implemented in many Asian, African, Mediterranean and Middle Eastern countries where G6PDD is common. Screening is associated with a reduced incidence of severe hyperbilirubinemia and kernicterus. In countries where G6PDD is historically less common, the increase in global population movement has raised the question as to whether G6PDD screening should be implemented throughout the world <sup>[31, 41]</sup>.

There was only one article about the relation between G6PD mutations, jaundice and its treatment in Iranian neonates; Zahedpasha et al <sup>[24]</sup>. Reported that there were no significant relationships between major mutations and icterus. A recent meta-analysis on five

papers represented that infants with G6PDD are about 4 times more at risk of hyperbilirubinemia compared with G6PD-normal infants <sup>[42]</sup>.

The current research also discussed kernicterus - a major complication of G6PDD. It is clear that G6PDD contributes to kernicterus via at least 2 mechanisms: firstly, severe hemolysis results in rising total bilirubin levels and subsequent accumulation of bilirubin in the brain. Secondly, G6PDD results in a reduced buffering capacity against bilirubin-induced reactive oxygen species <sup>[43, 44]</sup>. The second mechanism may explain why G6PD-deficient infants develop kernicterus at even at lower levels of total bilirubin. The risk of kernicterus in G6PD-deficient infants with total bilirubin serum levels above 20 mg/dL (342  $\mu$ mol/L) appears to be more severe than that associated with rhesus disease. Thus, in the presence of G6PDD, even more aggressive treatment of these infants is probably indicated <sup>[45-48]</sup>. The incidence of kernicterus in Iran has risen in recent years because of a variety of factors: firstly, newborns are often discharged from the hospital within 24 to 48 hours of birth, but total bilirubin levels often peak 4 to 5 days after birth. Secondly, the lack of proper monitoring at home allows the undiagnosed development of kernicterus <sup>[49-52]</sup>.

One limitation of this systematic review was the restricted number of studies evaluating complications of G6PDD. The current study suggests the planning of new screening studies, and follow-up of the G6PDD cases to record the frequency of jaundice and kernicterus. Another limitation was the high heterogeneities between the studies, despite analyzing only the population-based screening studies. However, because the individual articles were epidemiologic surveys, we would expect high heterogeneity <sup>[53]</sup>.

In conclusion, the prevalence of G6PDD in Iran is similar to most countries. Jaundice and kernicterus are the major complications of G6PDD. Therefore, it is necessary to provide good care for patients with G6PDD, and it is recommended for those hospitals to provide the result of G6PD testing as soon as possible after delivery, ideally before discharging newborn children.

## Acknowledgment

We are grateful to the Clinical Research Development Committee of Amirkola Children's Hospital and Student Research Committee of Babol University of Medical Sciences for their contribution to this study.



**Funding:** None.

**Ethical Approval:** This study obtained ethics committee of Babol University of Medical Sciences (IR.MUBABOL.REC.1398.052).

**Conflict of Interest:** None declared.

## References:

1. Howes RE, Piel FB, Patil AP, et al. G6PD deficiency prevalence and estimates of affected populations in malaria endemic countries: a geostatistical model-based map. *PLoS Med* 2012; 9(11): e1001339.
2. Domingo GJ, Satyagraha AW, Anvikar A, et al. G6PD testing in support of treatment and elimination of malaria: recommendations for evaluation of G6PD tests. *Malar J* 2013; 12: 391.
3. Bhutani VK, Zipursky A, Blencowe H, et al. Neonatal hyperbilirubinemia and Rhesus disease of the newborn: incidence and impairment estimates for 2010 at regional and global levels. *Pediatr Res* 2013; 74(Suppl 1): 86-100.
4. Leong A. Is there a need for neonatal screening of glucose-6-phosphate dehydrogenase deficiency in Canada? *McGill J Med* 2007; 10(1): 31-4.
5. Arese P, Gallo V, Pantaleo A, Turrini F. Life and death of glucose-6-phosphate dehydrogenase (G6PD) deficient erythrocytes—role of redox stress and band 3 modifications. *Transfus Med Hemother* 2012; 39(5): 328-34.
6. Pamba A, Richardson ND, Carter N, et al. Clinical spectrum and severity of hemolytic anemia in glucose 6-phosphate dehydrogenase-deficient children receiving dapsone. *Blood* 2012; 120(20): 4123-33.
7. Kaplan M, Hammerman C. Glucose-6-phosphate dehydrogenase deficiency and severe neonatal hyperbilirubinemia: a complexity of interactions between genes and environment. *Semin Fetal Neonatal Med* 2010; 15(3): 148-56.
8. Wusthoff CJ, Loe IM. Impact of bilirubin-induced neurologic dysfunction on neurodevelopmental outcomes. *Semin Fetal Neonatal Med*. 2015; 20(1): 52-7.
9. Maisels MJ. Managing the jaundiced newborn: a persistent challenge. *CMAJ*. 2015;187(5):335-43.
10. Cohan N, Karimi M, Khalili AH, Falahzadeh MH, Samadi B, Mahdavi RM. The efficacy of a neonatal screening programme in decreasing the hospitalization rate of patients with G6PD deficiency in southern Iran. *J Med Screen* 2010; 17(2): 66-7.
11. Arain YH, Bhutani VK. Prevention of Kernicterus in South Asia: role of neonatal G6PD deficiency and its identification. *Indian J Pediatr* 2014; 81(6): 599-607.
12. Kosaryan M, Nasehi MM, Karami H, et al. Neonatal screening for G6PD deficiency in Mazandaran Province, Iran 2007-2010. *Iran J Blood Cancer* 2011; 2(4): 113-6.
13. Moher D, Liberati A, Tetzlaff J, Altman DG, Group P. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med*. 2009;6(7):e1000097.
14. Available at: [http://www.crd.york.ac.uk/PROSPERO/display\\_record.php?ID=CRD42019119693](http://www.crd.york.ac.uk/PROSPERO/display_record.php?ID=CRD42019119693). Accessed on Feb 2, 2020.
15. Hoy D, Brooks P, Woolf A, et al. Assessing risk of bias in prevalence studies: modification of an existing tool and evidence of interrater agreement. *J Clin Epidemiol* 2012; 65(9): 934-9.
16. Khalesy N, Khosravi N, Haghighi M. Prevalence of Glucose-6-phosphate dehydrogenase deficiency in neonates born in Tehran-Iran (2008-09). *J Gorgan Uni Med Sci* 2012; 14(1): 100-5.
17. Abolghasemi H, Mehrani H, Amid A. An update on the prevalence of glucose-6-phosphate dehydrogenase deficiency and neonatal jaundice in Tehran neonates. *Clin Biochem* 2004; 37(3): 241-4.
18. Kazemi A, Norouzi H, Moghaddam MT, Naderloo A. Prevalence of Glucose-6-phosphate dehydrogenase (G6PD) deficiency of newborns in Rasol Akram and Ali Asghar hospitals of Tehran. *Iran South Med J* 2013; 16(1): 61-8 [Text in Persian].
19. Aletayeb SMH, Dehdashtian M, Aramesh MR, et al. Outcome of jaundice in neonates with ABO and Rh blood incompatibility and glucose-6-phosphate dehydrogenase deficiency. *Biomed Res* 2017; 28(8): 3440-4.
20. Boskabadi H, Omidian M, Mafinejad S. Prevalence and clinical manifestation of glucose-6-phosphate dehydrogenase deficiency in newborns with hyperbilirubinemia in Mashhad, Iran. *Maced J Med Sci* 2010; 3(4): 383-7.
21. Yousefi J, Mirzadeh M, Malek A. The role of hemolysis in inducing jaundice in the newborns with G6PD deficiency. *Iran J Pediatr* 2006;16(4):462-6.
22. Boskabadi H, Maamouri G, Mafinejad S, Rezagholizadeh F. Clinical course and prognosis of hemolytic jaundice in neonates in North East of Iran. *Maced J Med Sci* 2011; 4(4): 403-7.
23. Mahdavi MR, Kowsarian M, Roshan P, et al. Carrier frequency of three common G6PD gene mutations in

- neonates in province of Mazandaran, North of Iran, 2012. *Razi J Med Sci* 2014; 21(124): 64-70 [Text in Persian].
24. Zahedpasha Y, Kachouri MA, Niaki HA, Farhadi R. Comparison of molecular mutations of G6PD deficiency gene between icteric and nonicteric neonates. *Int J Mol Cell Med* 2013; 2(1): 14-20.
  25. Zahedpasha Y, Ahmadpour M, Niaki HA, Alaei E. Relation between Neonatal Icteric and Gilbert Syndrome in Glucose-6-Phosphate Dehydrogenase Deficient Subjects. *J Clin Diagn Res* 2014; 8(3): 63-5.
  26. Nock M, Johnson E, Krugman R, et al. Implementation and analysis of a pilot in-hospital newborn screening program for glucose-6-phosphate dehydrogenase deficiency in the United States. *J Perinatol* 2011; 31(2): 112-7.
  27. Goyal M, Garg A, Goyal MB, et al. Newborn screening for G6PD deficiency: A 2-year data from North India. *Indian J Public Health* 2015; 59(2): 145-8.
  28. Thielemans L, Gornawun G, Hanboonkunupakarn B, et al. Diagnostic performances of the fluorescent spot test for G6PD deficiency in newborns along the Thailand-Myanmar border: A cohort study. *Wellcome open Res* 2018; 3: 1.
  29. Zhang J, Cui Y, Wang X, et al. Prevalence of Glucose-6-Phosphate Dehydrogenase Deficiency in Sichuan, China. *Clin Lab* 2018; 64(3): 383-6.
  30. Castro S, Weber R, Dadalt V, et al. Prevalence of G6PD deficiency in newborns in the south of Brazil. *J Med Screen* 2006; 13(2): 85-6.
  31. Elalla SA, Tawfik M, Barseem N, Moustafa W. Prevalence of glucose-6-phosphate dehydrogenase deficiency in neonates in Egypt. *Ann Saudi Med* 2017; 37(5): 362-5.
  32. Guellouz N, Ben IM, Ouederni M, et al. Neonatal screening of G6PD deficiency in Tunisia. *Arch Inst Pasteur Tunis* 2010; 87(1-2): 69-76.
  33. Muzaffer MA. Neonatal screening of glucose-6-phosphate dehydrogenase deficiency in Yanbu, Saudi Arabia. *J Med Screen* 2005; 12(4): 170-1.
  34. Moosazadeh M, Nekoei-Moghadam M, ALIRAM-ZANY M, Amiresmaili M. Identification of mutation of Glucose-6-Phosphate Dehydrogenase (G6PD) in Iran: Meta-analysis study. *Iran J Public Health* 2013; 42(9): 1007-15.
  35. Luzzatto L, Arese P. Favism and glucose-6-phosphate dehydrogenase deficiency. *New England J Med* 2018; 378(1): 60-71.
  36. Laosombat V, Sattayasevana B, Chotsampancharoen T, Wongchanchailert M. Glucose-6-phosphate dehydrogenase variants associated with favism in Thai children. *Int J Hematol* 2006; 83(2): 139-43.
  37. Vulliamy T, D'urso M, Battistuzzi G, et al. Diverse point mutations in the human glucose-6-phosphate dehydrogenase gene cause enzyme deficiency and mild or severe hemolytic anemia. *Proc Natl Acad Sci U S A* 1988; 85(14): 5171-5.
  38. Rahimi Z, Vaisi-Raygani A, Nagel RL, Muniz A. Molecular characterization of glucose-6-phosphate dehydrogenase deficiency in the Kurdish population of Western Iran. *Blood Cells Mol Dis* 2006; 37(2): 91-4.
  39. Cappellini MD, Fiorelli G. Glucose-6-phosphate dehydrogenase deficiency. *Lancet* 2008; 371(9606): 64-74.
  40. Chien Y-H, Lee N-C, Wu S-T, et al. Changes in incidence and sex ratio of glucose-6-phosphate dehydrogenase deficiency by population drift in Taiwan. *Southeast Asian J Trop Med Public Health* 2008; 39(1): 154-61.
  41. Watchko J, Kaplan M, Stark A, et al. Should we screen newborns for glucose-6-phosphate dehydrogenase deficiency in the United States? *J Perinatol* 2013; 33(7): 499-504.
  42. Liu H, Liu W, Tang X, Wang T. Association between G6PD deficiency and hyperbilirubinemia in neonates: a meta-analysis. *Pediatr Hematol Oncol* 2015; 32(2): 92-8.
  43. de Gurrola GC, Araúz JJ, Durán E, et al. Kernicterus by glucose-6-phosphate dehydrogenase deficiency: a case report and review of the literature. *J Med Case Reports* 2008; 2: 146.
  44. Kaplan M, Hammerman C. Glucose-6-phosphate dehydrogenase deficiency: a hidden risk for kernicterus. *Semin Perinatol* 2004; 28(5): 356-64.
  45. Slusher TM, Vreman HJ, McLaren DW, et al. Glucose-6-phosphate dehydrogenase deficiency and carboxyhemoglobin concentrations associated with bilirubin-related morbidity and death in Nigerian infants. *J Pediatr* 1995; 126(1): 102-8.
  46. Brown WR, Boon WH. Hyperbilirubinemia and kernicterus in glucose-6-phosphate dehydrogenase-deficient infants in Singapore. *Pediatrics* 1968; 41(6): 1055-62.
  47. Gibbs WN, Gray R, Lowry M. Glucose-6-phosphate dehydrogenase deficiency and neonatal jaundice in Jamaica. *Br J Haematol* 1979; 43(2): 263-74.
  48. Hannon PR, Willis SK, Scrimshaw SC. Persistence of maternal concerns surrounding neonatal jaundice: an

- exploratory study. *Arch Pediatr Adolesc Med* 2001; 155(12): 1357-63.
49. Olusanya BO, Ogunlesi TA, Slusher TM. Why is kernicterus still a major cause of death and disability in low-income and middle-income countries? *Arch Dis Child* 2014; 99(12): archdischild-2013-305506.
  50. Watchko JF. Identification of neonates at risk for hazardous hyperbilirubinemia: emerging clinical insights. *Pediatr Clin North Am* 2009; 56(3): 671-87.
  51. Ives K. Preventing kernicterus: a wake-up call. *Arch Dis Child Fetal Neonatal Ed* 2007; 92(5): F330-F1.
  52. Bhutani V, Johnson L. Kernicterus in the 21st century: frequently asked questions. *J Perinatol* 2009; 29(Suppl 1): S20-4.
  53. Zamani M, Derakhshan M, Zamani V, Shokri-Shirvani J. The prevalence of *Helicobacter pylori* infection worldwide knowns and unknowns. Authors' reply. *Aliment Pharmacol Ther* 2018; 47(9): 1331-2.
  54. ZahedPasha Y. The prevalence of G6PD deficiency in the newborns of Babol, 1995. *J Babol Univ Med Sci* 1999; 1(4): 19-25 [Text in Persian].
  55. Movahhed A, Farrokhi S. Incidence Rate of G6PD Deficiency in Newborns in Bushehr/Iran. *Iran J Pediatr* 2003; 13(1): 55-8 [Text in Persian].
  56. Daliri S, Asadollahi K, Rahimi N, Sayehmiri K. Incidence of glucose-6-phosphate dehydrogenase deficiency in malaria-prone regions of Fars province. *Tehran Univ Med J* 2017; 75(9): 669-74 [Text in Persian].
  57. Iranpour R, Hashemipour M, Talaei S-M, et al. Newborn screening for glucose-6-phosphate dehydrogenase deficiency in Isfahan, Iran: a quantitative assay. *J Med Screen* 2008; 15(2): 62-4.
  58. Mohammadzadeh A, Jafarzadeh M, Farhat AS, et al. Prevalence of glucose-6-phosphate dehydrogenase deficiency in neonates of Northeast of Iran. *J Clin Med Res* 2009; 4(8).
  59. Kosaryan M, Mahdavi M, Jalali H, Roshan P. Comparison of fluorescent spot test, decolorization test and quantitative enzyme assay in detection of G6PD enzyme deficiency. *J Gorgan Uni Med Sci*. 2015;17(3): 108-13 [Text in Persian].
  60. Alidalaki S, Negahban BT, Houlakouei M, et al. Investigation of glucose-6-phosphate dehydrogenase deficiency in Rafsanjan, Autumn 2004. *J Rafsanjan Univ Med Sci* 2008 [Text in Persian].
  61. Khalili D, Jafroodi M, Sajedi S, et al. Survey of the prevalence of glucose-6-phosphate dehydrogenase deficiency in Rasht-Iran. *J Guilan Uni Med Sci* 2007; 16(63): 51-6 [Text in Persian].
  62. Keihanian F, Basirjafari S, Darbandi B, Saeidinia A, Jafroodi M, Sharafi R, et al. Comparison of quantitative and qualitative tests for glucose-6-phosphate dehydrogenase deficiency in the neonatal period. *Int J Lab Hematol* 2017; 39(3): 251-60.
  63. Kosaryan M, Mahdavi MR, Jalali H, Roshan P. Why does the Iranian national program of screening newborns for G6PD enzyme deficiency miss a large number of affected infants? *Pediatr Hematol Oncol*. 2014;31(1):95-100.
  64. Nazari H, Habibeh Najar H, Emadi A, et al. Prevalence of G6PD deficiency in neonates referred to Semnan University of Medical Science's screening Lab. *Medical Laboratory Journal*. 2011;5(2):66-70 [Text in Persian].
  65. Norbakhsh Smk, Kasiri k-A, Jalil A, Hashemzade m. Determining the frequency of glucose-6-phosphate dehydrogenase deficiency in newborn infants in Shahrekord. *J of Shahrekord Uuniversity of Medical Sciences*. 2013;15(5):30-7 [Text in Persian].
  66. Ansari-Moghaddam A, Adineh HA, Mohammadi M, et al. Prevalence of glucose-6-phosphate dehydrogenase (G6PD) deficiency in malaria endemic region of Iran (Sistan and Baluchestan Province): Epidemiological profile and trends over time. *Asian Pac J Trop Dis* 2017; 7(10): 587-91.